



Neff constraints on the portal interactions with hidden sector

arxiv:2111.xxxxx

-Pranjal Ralegankar
University of Illinois at Urbana-Champaign


Collaborators: Peter Adshead and
Jessie Shelton

Outline

- Background and motivation
- Physics behind Neff constraints
 - Gauged B-L with right handed neutrinos
- Extending to more general hidden sectors
- Summary



Background and Motivation

- All known particles are charged under Standard Model gauge symmetries, i.e. they interact with Strong and Electroweak forces.




Background and Motivation

- But there is no fundamental requirement for all particles in the universe to be charged under Standard Model symmetries.




Background and Motivation

- Any such new particle can have renormalizable portal couplings with the Standard Model particles that obey the Standard Model gauge symmetries.




Background and Motivation

- Collider experiments can probe portal interactions




Background and Motivation

- So can Cosmic microwave background (CMB) spectrum probe portal interactions



Background and Motivation


- So can Cosmic microwave background (CMB) spectrum probe portal interactions



Physics behind Neff constraints: B-L example

$$L_{int} \supset -\frac{1}{4} F'_{\mu\nu} F^{\mu\nu'} + g' Z'_\mu J_{B-L,SM}^\mu - g' Z'_\mu \sum_i \bar{\nu}_{R,i} \gamma^\mu \nu_{R,i} + \frac{1}{2} M_{Z'}^2 Z'^\mu Z'_\mu$$


↑
Standard Model B-L current ↑
Right handed neutrinos →
Stueckelberg mass



Physics behind Neff constraints: Boltzmann equations

$$L_{int} \supset -\frac{1}{4} F'_{\mu\nu} F^{\mu\nu'} + g' Z'_\mu J_{B-L,SM}^\mu - g' Z'_\mu \sum_i \bar{\nu}_{R,i} \gamma^\mu \nu_{R,i} + \frac{1}{2} M_{Z'}^2 Z'^\mu Z'_\mu$$

↑
Standard Model B-L current ↑
Right handed neutrinos →
Stueckelberg mass



Boltzmann equations:

$$\frac{d\rho_{SM}}{dt} + 3H(1+w_{SM})\rho_{SM} = -C$$

$$\frac{d\rho_{\nu_R}}{dt} + 4H\rho_{\nu_R} = C$$

$$H = \frac{\sqrt{\rho_{SM} + \rho_{\nu_R}}}{\sqrt{3}M_{pl}}$$

$$C = \frac{1}{32\pi^4} \sum_f \int ds (s - 4m_f^2) s \sigma_{ff \rightarrow \nu_R \nu_R} [T_{SM} G(\sqrt{s}/T_{SM}) - T_{\nu_R} G(\sqrt{s}/T_{\nu_R})]$$

↑
Energy transfer collision term

Physics behind Neff constraints: Energy density evolution without thermalization



Boltzmann equations:

$$\frac{d\rho_{SM}}{dt} + 3H(1+w_{SM})\rho_{SM} = -C$$

$$\frac{d\rho_{\nu_R}}{dt} + 4H\rho_{\nu_R} = C$$

$$H = \frac{\sqrt{\rho_{SM} + \rho_{\nu_R}}}{\sqrt{3}M_{pl}}$$

$$C = \frac{1}{32\pi^4} \sum_f \int ds (s - 4m_f^2) s \sigma_{ff \rightarrow \nu_R \nu_R} [T_{SM} G(\sqrt{s}/T_{SM}) - T_{\nu_R} G(\sqrt{s}/T_{\nu_R})]$$



If ν_R is always in equilibrium with the Standard Model plasma

Only the forward part of the collision term

Physics behind Neff constraints: Energy density evolution without thermalization

Boltzmann equations:



$$\frac{d\rho_{SM}}{dt} + 3H(1+w_{SM})\rho_{SM} = -C$$

$$\frac{d\rho_{\nu_R}}{dt} + 4H\rho_{\nu_R} = C$$

$$H = \frac{\sqrt{\rho_{SM} + \rho_{\nu_R}}}{\sqrt{3}M_{pl}}$$

$$C = \frac{1}{32\pi^4} \sum_f \int ds (s - 4m_f^2) s \sigma_{ff \rightarrow \nu_R \nu_R} [T_{SM} G(\sqrt{s}/T_{SM}) - T_{\nu_R} G(\sqrt{s}/T_{\nu_R})]$$

Final energy density proportional to the strength of the portal interaction



If ν_R is always in equilibrium with the Standard Model plasma

Only the forward part of the collision term

Physics behind Neff constraints: Energy density evolution



Boltzmann equations:

$$\frac{d\rho_{SM}}{dt} + 3H(1+w_{SM})\rho_{SM} = -C$$

$$\frac{d\rho_{\nu_R}}{dt} + 4H\rho_{\nu_R} = C$$

$$H = \frac{\sqrt{\rho_{SM} + \rho_{\nu_R}}}{\sqrt{3}M_{pl}}$$

$$C = \frac{1}{32\pi^4} \sum_f \int ds (s - 4m_f^2) s \sigma_{ff \rightarrow \nu_R \nu_R} [T_{SM} G(\sqrt{s}/T_{SM}) - T_{\nu_R} G(\sqrt{s}/T_{\nu_R})]$$



If ν_R is always in equilibrium with the Standard Model plasma

Only the forward part of the collision term

Physics behind Neff constraints: Energy density evolution with thermalization

Boltzmann equations:

$$\frac{d\rho_{SM}}{dt} + 3H(1 + w_{SM})\rho_{SM} = -C$$

$$\frac{d\rho_{\nu_R}}{dt} + 4H\rho_{\nu_R} = C$$



$$H = \frac{\sqrt{\rho_{SM} + \rho_{\nu_R}}}{\sqrt{3}M_{pl}}$$

$$C = \frac{1}{32\pi^4} \sum_f \int ds (s - 4m_f^2) s \sigma_{ff \rightarrow \nu_R \nu_R} [T_{SM} G(\sqrt{s}/T_{SM}) - T_{\nu_R} G(\sqrt{s}/T_{\nu_R})]$$

Final energy density largely insensitive to the strength of the portal interaction



$$\frac{\rho_{\nu_R}}{\rho_{SM}} \propto \frac{g_{\nu_R}}{g_{*SM}}$$

Degrees of freedom





If ν_R is always in equilibrium with the Standard Model plasma


Physics behind Neff constraints: Translating to constraints




Physics behind Neff constraints: Translating to constraints




Physics behind Neff constraints: Most relevant when thermally decoupled




Comparing B-L constraints: Future constraint will extend to much larger parameter space



Comparing B-L constraints: Neff constraints dominant for $M_{Z'} > 0.1$ MeV




Comparing B-L constraints: Improving over previous analysis




1. K. N. Abazajian and J. Heeck, Phys. Rev. D 100 (2019) 075027.


Extending to general hidden sector



Extending to general hidden sector




Extending to general hidden sector



Neff constraints on the portal interactions with the minimal HS act as conservative constraints on extended HS.


Leaked energy independent of details within HS



$$\frac{\rho_{HS}}{\rho_{SM}} \propto \frac{1}{g_{*SM}(M_{Z'}/8)} \frac{M_{pl}}{M_{Z'}} L$$

$$L = M_{Z'} \int ds \frac{s - 4m_f^2}{s\sqrt{s}} \sigma_{ff \rightarrow \nu_R \nu_R} \propto g'^2$$

↑
Leak factor



Leaked energy independent of details within HS



Independent of degrees of Freedom or the temperature In HS!



$$\frac{\rho_{HS}}{\rho_{SM}} \propto \frac{1}{g_{*SM}(M_{Z'}/8)} \frac{M_{pl}}{M_{Z'}} L$$

$$L = M_{Z'} \int ds \frac{s - 4m_f^2}{s\sqrt{s}} \sigma_{ff \rightarrow \nu_R \nu_R} \propto g'^2$$


Leak factor



Leaked energy independent of details within HS: Depends only on one BSM coupling

Beyond Standard Model


Independent of degrees of Freedom or the temperature In HS!



$$\frac{\rho_{HS}}{\rho_{SM}} \propto \frac{1}{g_{*SM}(M_{Z'}/8)} \frac{M_{pl}}{M_{Z'}} L$$

$$L = M_{Z'} \int ds \frac{s - 4m_f^2}{s\sqrt{s}} \sigma_{ff \rightarrow \nu_R \nu_R} \propto g'^2$$

Leak factor




Leaked energy independent of details within HS: Depends only on one BSM coupling

Independent of degrees of Freedom or the temperature In HS!



$$\frac{\rho_{HS}}{\rho_{SM}} \propto \frac{1}{g_{*SM}(M_{Z'}/8)} \frac{M_{pl}}{M_{Z'}} L$$

$$L = M_{Z'} \int ds \frac{s - 4m_f^2}{s\sqrt{s}} \sigma_{ff \rightarrow \nu_R \nu_R} \propto g'^2$$


Leak factor



Neff constraints applicable for wide class of hidden sectors




Reinterpreting the utility of Neff: Typically discussed as constraint on decoupling temperature




arxiv:1903:04763

Reinterpreting the utility of Neff: Instead think of as constraint on a wide class of hidden sectors





Decoupling temperature of new light particle



Summary

- N_{eff} constraints are most relevant when the relativistic particles remain out-of-equilibrium with the Standard Model plasma
- As N_{eff} measurements improve, the thermalization threshold will be pushed further away
- N_{eff} measurements can provide conservative constraints on the portal couplings that are applicable for a large class of thermally decoupled hidden sectors.




Backup slides

Physics behind Neff constraints: Boltzmann equations

$$L_{int} \supset -\frac{1}{4} F'_{\mu\nu} F^{\mu\nu'} + g' Z'_\mu J_{B-L,SM}^\mu - g' Z'_\mu \sum_i \bar{\nu}_{R,i} \gamma^\mu \nu_{R,i} + \frac{1}{2} M_{Z'}^2 Z'^\mu Z'_\mu$$

↑
Standard Model B-L current ↑
Right handed neutrinos →
Stueckelberg mass



Boltzmann equations:

$$\frac{d\rho_{SM}}{dt} + 3H(1+w_{SM})\rho_{SM} = -C$$

$$\frac{d\rho_{\nu_R}}{dt} + 4H\rho_{\nu_R} = C$$


$$H = \frac{\sqrt{\rho_{SM} + \rho_{\nu_R}}}{\sqrt{3}M_{pl}}$$

$$C = \frac{1}{32\pi^4} \sum_f \int ds (s - 4m_f^2) s \sigma_{ff \rightarrow \nu_R \nu_R} [T_{SM} G(\sqrt{s}/T_{SM}) - T_{\nu_R} G(\sqrt{s}/T_{\nu_R})]$$

Physics behind Neff constraints: Boltzmann equations

$$L_{int} \supset -\frac{1}{4} F'_{\mu\nu} F^{\mu\nu'} + g' Z'_\mu J_{B-L,SM}^\mu - g' Z'_\mu \sum_i \bar{\nu}_{R,i} \gamma^\mu \nu_{R,i} + \frac{1}{2} M_{Z'}^2 Z'^\mu Z'_\mu$$

↑
Standard Model B-L current
↑
Right handed neutrinos
→ Stueckelberg mass



Boltzmann equations:

$$\frac{d\rho_{SM}}{dt} + 3H(1+w_{SM})\rho_{SM} = -C$$

$$\frac{d\rho_{\nu_R}}{dt} + 4H\rho_{\nu_R} = C$$

$$H = \frac{\sqrt{\rho_{SM} + \rho_{\nu_R}}}{\sqrt{3}M_{pl}}$$


$$C = \frac{1}{32\pi^4} \sum_f \int ds (s - 4m_f^2) s \sigma_{ff \rightarrow \nu_R \nu_R} [T_{SM} G(\sqrt{s}/T_{SM}) - T_{\nu_R} G(\sqrt{s}/T_{\nu_R})]$$

↑
Energy transfer collision term

Physics behind Neff constraints: Boltzmann equations

$$L_{int} \supset -\frac{1}{4} F'_{\mu\nu} F^{\mu\nu'} + g' Z'_\mu J_{B-L,SM}^\mu - g' Z'_\mu \sum_i \bar{\nu}_{R,i} \gamma^\mu \nu_{R,i} + \frac{1}{2} M_{Z'}^2 Z'^\mu Z'_\mu$$

↑
Standard Model B-L current ↑
Right handed neutrinos →
Stueckelberg mass



Boltzmann equations:

$$\frac{d\rho_{SM}}{dt} + 3H(1+w_{SM})\rho_{SM} = -C$$

$$\frac{d\rho_{\nu_R}}{dt} + 4H\rho_{\nu_R} = C$$

$$H = \frac{\sqrt{\rho_{SM} + \rho_{\nu_R}}}{\sqrt{3}M_{pl}}$$

Integral over Mandelstam s (or centre of mass energy)


$$C = \frac{1}{32\pi^4} \sum_f \int ds (s - 4m_f^2) s \sigma_{ff \rightarrow \nu_R \nu_R} [T_{SM} G(\sqrt{s}/T_{SM}) - T_{\nu_R} G(\sqrt{s}/T_{\nu_R})]$$

↑
Energy transfer collision term

Physics behind Neff constraints: Boltzmann equations

$$L_{int} \supset -\frac{1}{4} F'_{\mu\nu} F^{\mu\nu'} + g' Z'_\mu J_{B-L,SM}^\mu - g' Z'_\mu \sum_i \bar{\nu}_{R,i} \gamma^\mu \nu_{R,i} + \frac{1}{2} M_{Z'}^2 Z'^\mu Z'_\mu$$

↑
Standard Model B-L current ↑
Right handed neutrinos →
Stueckelberg mass



Boltzmann equations:

$$\frac{d\rho_{SM}}{dt} + 3H(1+w_{SM})\rho_{SM} = -C$$

$$\frac{d\rho_{\nu_R}}{dt} + 4H\rho_{\nu_R} = C$$

$$H = \frac{\sqrt{\rho_{SM} + \rho_{\nu_R}}}{\sqrt{3}M_{pl}}$$

CM frame cross-section


$$C = \frac{1}{32\pi^4} \sum_f \int ds (s - 4m_f^2) s \sigma_{f\bar{f} \rightarrow \nu_R \bar{\nu}_R} [T_{SM} G(\sqrt{s}/T_{SM}) - T_{\nu_R} G(\sqrt{s}/T_{\nu_R})]$$

↑
Energy transfer collision term

Physics behind Neff constraints: Boltzmann equations

$$L_{int} \supset -\frac{1}{4} F'_{\mu\nu} F^{\mu\nu'} + g' Z'_\mu J_{B-L,SM}^\mu - g' Z'_\mu \sum_i \bar{\nu}_{R,i} \gamma^\mu \nu_{R,i} + \frac{1}{2} M_{Z'}^2 Z'^\mu Z'_\mu$$

↑ Standard Model B-L current
 ↑ Right handed neutrinos
 → Stueckelberg mass



Boltzmann equations:


$$\frac{d\rho_{SM}}{dt} + 3H(1+w_{SM})\rho_{SM} = -C$$

$$\frac{d\rho_{\nu_R}}{dt} + 4H\rho_{\nu_R} = C$$

$$H = \frac{\sqrt{\rho_{SM} + \rho_{\nu_R}}}{\sqrt{3}M_{pl}}$$

$$C = \frac{1}{32\pi^4} \sum_f \int ds (s - 4m_f^2) s \sigma_{ff \rightarrow \nu_R \nu_R} [T_{SM} G(\sqrt{s}/T_{SM}) - T_{\nu_R} G(\sqrt{s}/T_{\nu_R})]$$

↑
Energy transfer collision term



Phase-space factor encoding Fermi-Dirac distribution